



garlic

APPLICATION TIMES, MODES AND DOSE

1st foliar application: at the appearance of the first real leaves.

Dose: 1,5-2 L/Ha.

2nd foliar application: 30 days after the 1st application, at least 3 weeks before harvesting.

Dose: 1,5-2 L/Ha.

BENEFITS AND ADVANTAGES

Given that garlic has a “centre of high metabolic activity”, it is the perfect crop on which to use **BROTOMAX**[®], due to its influence on secondary metabolism.

- Greener leaves. More chlorophyll.
- Improves the plant's bearing. More vigorous plants.
- Does not soften the clove.
- Provides quality: better skin and a higher calibre.
- Greater yield at harvest.
- Helps the plant to overcome stress produced during cultivation because of fungicides and herbicides treatments.
- Increases the level of polyphenols, resulting in an increase in antioxidant potential.

SEEDS

Use **BROTOMAX**[®] mixed with fungicides.

Dose: 1,5 litres per 1000kg of shelled cloves.

Advantages:

- Improves emergence (less losses when planting).
- Favours the growth of the root.

RECOMMENDATIONS

Use of **CORRIZ•AM**[®] (root application) is recommended at plantation to cause greater proliferation of absorbent hairs which in turn results in better plants setting.

Dose (irrigation application): 5 L/Ha.