



broccoli and cauliflower

APPLICATION TIMES, MODES AND DOSE

1st application: in the first localised irrigation.

Dose: 5 L/Ha.

2nd foliar application: when the cluster (head) begins to form, which is usually a month after the 1st application (planting). This is when the plant needs nitrogen the most. Compatible with other products that can be applied at the same time.

Dose: 3 L/Ha.

CORRIZ•AM® with **BROTOMAX®**, is recommended via irrigation, in the first application.

Dose: 5 L/Ha.

BENEFITS AND ADVANTAGES

- Greater root development and better ground setting.
- Vigorous plants.
- Weight increase.
- Foliar application increases the size of the leaves, ensuring full coverage.
- Greater resistance to rot and neck diseases (*Sclerotinia*, *Pythium*...).
- Better storage preservation, obtaining more vigorous plants, resistant to bacterial soft rots and to aqueous softening caused by fungi.

RECOMMENDATIONS

- An application of **BROTOMAX®**, in prolonged water logging due to heavy rains is recommended as it is compatible with copper and specific fungicides and so dramatically helps the recovery of surviving plants.

Dose: 3 L/Ha.

- Once the stalk is formed, the application of **QUASSAR®** is recommended to prevent the formation of dew, thus better dryness and as a result prevents subsequent rot.

Dose: 0,15-0,25%.

- The use of **BIOHIDRAMAR® CREMA** would boost plant protection products.

Foliar dose: 2 L/Ha.