



avocado

APPLICATION TIMES, MODES AND DOSE

1st foliar application: at winter budding.

2nd foliar application: during maturation.

3rd foliar application: during post-harvest.

Dose:

foliar application: 3 L/Ha.

irrigation application: 5-6 L/Ha.

BENEFITS AND ADVANTAGES

- Increase in the number of buds sprouted.
- Ensures good stimulation of sprouting buds. Optimum balance between vegetation (foliar surface) and number of flowers, better lining of the tree.
- Strengthens self-defence mechanisms of the tree to overcome attacks of *Phytophthora* spp.
- Uniformity of fruit and more homogeneous maturation.
- Better quality and consistency of fruit skin.
- Increases the size and uniformity of fruit calibre.
- Better conservation of the fruit.
- With post-harvest treatment, an increase in reserves for the next season is guaranteed.

RECOMMENDATIONS

Apply **BIOHIDRAMAR® CUAJE** from the beginning of flowering until petals are visible thus boosting the fruit-bearing potential of the crop besides increasing pollen quality and the number of uniform and fertile flowers as well, thus insuring better fruit setting.

Dose (foliar application): 0,1% (1 L/1000L).

Several applications of **AM•EN®** from marble-sized fruit to egg-sized (approximately every 15-21 days) is recommended in order to boost fruit swelling. Besides, boosting fruit gripping, greater calibre homogeneity and improvement in fruit quality.

Dose (foliar application): 0,3% (3 L/1000 L).

We recommend using **LANZADERA® CALCIO** to increase the quality and the shelf life of the fruit.

Dose:

foliar application: 0,3% (3 L/1000 L) or

irrigation application: 5 L/Ha.

Applications of **LANZADERA® POTASIO** is recommended to encourage fruit maturation and swelling.

Dose:

foliar application: 0,25% (2,5 L/1000 L) or

irrigation application: 5 L/Ha.

A mixture with **QUASSAR®** is recommended to heighten the effectiveness of phytosanitary and nutritional treatments due to its high surfactant, adherent and penetrating power.

Foliar dose: 0,2-0,3%.

When planting begins and at times of stress **CORRIZ•AM®** is recommended to achieve:

- Proliferation of absorbent hairs and a larger root mass.
- Generating vegetative improvement and a larger trunk diameter.
- Helps to overcome moments caused by biotic and abiotic agents.
- Promotes the multiplication and lengthening of the roots.
- Stimulates mycorrhization.

Dose (irrigation application): 5-6 L/Ha.



BROTO MAX®